Quadratic Forms and Symmetric Bilinear Functions

LA3 \diamond **1.** Find the matrix of a bilinear function in new basis (e'_1, e'_2, e'_3) , if it has the matrix

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

in the old basis (e_1, e_2, e_3) , where

$$e'_1 = e_1 - e_2$$
, $e'_2 = e_1 + e_3$, $e'_3 = e_1 + e_2 + e_3$.

LA32. Prove that the determinant of a skew-symmetric integral matrix is a square of some integer number.

LA3 \diamond **3.** Suppose f is a skew-symmetric bilinear function on a space V, W is a subspace of V and W^{\perp} is its orthogonal complement with respect to f. Prove that dim W-dim($W \cap W^{\perp}$) is an even number.

LA304. Are the bilinear functions with matrices

$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & -1 \\ 3 & -1 & 3 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 3 & 0 \\ 3 & 1 & 1 \\ 0 & 1 & 5 \end{pmatrix}$$

isomorphic to each other?

LA3 \diamond **5.** Find all $\lambda \in \mathbb{R}$ for which the quadratic form

$$q(x) = 5x_1^2 + x_2^2 + \lambda x_3^2 + 4x_1x_2 - 2x_1x_3 - 2x_2x_3$$

is positive definite.

LA3 \diamond **6.** Find the positive and negative inertial indexes of the quadratic form $q(x) = \operatorname{tr} x^2$ on the space $\operatorname{Mat}_n(\mathbb{R})$.